

The Lowest Total Irregularity of a Completely Segregated ∞ –Bicyclic Graph

Mahadevaswamy B.S.

Assistant Professor, Department of Mathematics
Maharanis science college for women,Mysore-570005, India

<p style="text-align: center;">Article History</p> <p style="text-align: center;">Article Submission</p> <p style="text-align: center;">16 February 2016</p> <p style="text-align: center;">Revised Submission</p> <p style="text-align: center;">29 April 2016</p> <p style="text-align: center;">Article Accepted</p> <p style="text-align: center;">18 May 2016</p> <p style="text-align: center;">Article Published</p> <p style="text-align: center;">30 June 2016</p>	<p style="text-align: center;">Abstract</p> <p>A simple linked graph with exactly the same number of edges as vertices plus one is called a bicyclic graph. A completely segregated bicyclic network is one in which any two neighboring vertices have different degrees. Total Irregularity of a graph is defined as: $irr_t(G) = \frac{1}{2} \sum_{u,v \in V(G)} d_G(u) - d_G(v)$. In this paper, total irregularity of totally segregated ∞- bicyclic graph is discussed and some properties of totally segregated ∞- bicyclic graph G with $\Delta=4$ and $n_4(G)=1$ is found. The basic bicycle denoted by $\infty(p,q,1)$ is obtained from two vertex-disjoint cycles C_p and C_q by identifying one vertex of C_p and one vertex of C_q.</p> <p>Keywords: Total irregularity, Totally segregated ∞-bicyclic graph, Basic bicycle $\infty(p,q,1)$, Degree sequence.</p>
---	---

1. Introduction

In this paper we consider only simple undirected connected graphs. Let $G=(V,E)$ be a graph of order n and size m . For $u, v \in V(G)$, we denote the number of edges incident to v by $\deg_G(v)$ or $d_G(v)$. The length of the shortest path from u to v by $d(u, v)$. $N_G(v)=\{w:d(v,w)=1\}$ denotes neighbourhood of v . For a graph G , let $\Delta(G)$ be the maximum degree of the vertices of G . Degree sequence of a graph is a non-increasing sequence of the vertex degrees of the graph.

One among them is the total irregularity of a graph introduced by Abdo, Brandt and Dimitrov [1], which is defined as

$$irr_t(G) = \frac{1}{2} \sum_{u,v \in V(G)} |d_G(u) - d_G(v)|. \quad \dots\dots\dots(1)$$

Graphs that have unique degrees for any two neighboring vertices are referred to as completely segregated, and this is what Jackson and Entringer [9] examined. In their study of a particular situation, Jorry T. F. and Parvathy K. S. [10] looked at trees where the degrees of any two neighboring

vertices varied by a constant $k \neq 0$, and these graphs are named as k -segregated tree. A graph G is k -segregated, if $|d_G(u) - d_G(v)| = k \neq 0$ for all edges $uv \in E(G)$ [10]. However, in many applications and problems, it is of great importance to know how irregular a given graph is. In [1] the graphs with maximal total irregularity are determined and the upper bound of the set $\{irr_t(G) : |V(G)| = n\}$ is obtained as:

$$irr_t(G_{max}) = \frac{1}{12} (2n^3 - 3n^2 - 2n), \text{ if } n \text{ even.}$$

$$= \frac{1}{12} (2n^3 - 3n^2 - 2n + 3), \text{ if } n \text{ odd.}$$

In [1] authors also showed that, If T is a tree on n vertices, $irr_t(T) \leq (n-1)(n-2)$, and the star $K_{1,n-1}$ has the maximal total irregularity among all trees of same order. Star on n vertices $K_{1,n-1}$, $n \geq 3$ is totally segregated tree (TST), particularly it is $(n-2)$ -segregated tree. Hence, maximum total irregularity of totally segregated trees on n vertices is $(n-1)(n-2)$, and the maximum is attained for $(n-2)$ -segregated tree i.e. for $K_{1,n-1}$. Characterization of totally segregated trees with $\Delta = 3$ and minimum total irregularity of totally segregated trees on n vertices is found in [11].

Theorem [11]

Let T be a TST on n vertices and $\Delta(T) = 3$. Then $n_3(T) - 1 \leq n_2(T) \leq 2n_3(T) + 1$.

In [2], they characterized the non-regular graphs with minimal total irregularity and thereby resolved the conjecture by Zhu, You and Yang [12] about the lower bound on the minimal total irregularity of non-regular connected graphs. Abdo and Dimitrov [2] also characterized the non-regular graphs with the second and the third smallest total irregularity.

Throughout this paper let n_i denote the number of vertices with degree i . Here, equation (1) is rewritten as:

$$irr_t(G) = \sum_{d_i > d_j} (d_i - d_j)(n_{d_i} - n_{d_j}) \dots\dots\dots(2)$$

where d_i and d_j denote different vertex degrees in G . Keep in mind that a graph's degree sequence determines how erratic it is overall. The total irregularity of graphs with degree sequences that are equal is the same.

In this paper we find minimum total irregularity of totally segregated ∞ -bicyclic graph on n vertices. For the sake of convenience totally segregated bicyclic graph is called a TSB-graph. We represent degree sequence of graph in the following form $(k^{n_k}, (k-1)^{n_{k-1}}, \dots, 3^{n_3}, 2^{n_2}, 1^{n_1})$.

2. Totally Segregated Bicyclic (TSB) graphs on n vertices with minimum Total Irregularity:

In The writers of [13] provide many bicyclic graph types. A simple linked graph with exactly the same number of edges as vertices plus one is called a bicyclic graph. The basic bicycle, denoted by $\infty(p, q, 1)$, is obtained from two vertex-disjoint cycles C_p and C_q by identifying one vertex of C_p and one vertex of C_q , where $p, q \geq 3$. See Figure 1.




Figure 1

A totally segregated bicyclic (TSB) graph is a bicyclic graph which is totally segregated.

See Figure 2.




Figure 2

In this section, we determine minimum total irregularity of totally segregated ∞ -bicyclic graph on n vertices. Observe that any ∞ -bicyclic graph G is obtained from a basic bicycle ∞ -($p, q, 1$) by attaching trees to some of its vertices. If G is obtained from ∞ -($p, q, 1$) by attaching trees to some of its vertices then we call G as bicyclic graph with basic bicycle ∞ -($p, q, 1$). Obviously B_n is the set of those graphs each of which is a bicyclic graph with basic bicycle ∞ -($p, q, 1$), ($p \geq 3, q \geq 3$) which is called ∞ -bicyclic graph for convenience.

Remark 2.1

For $n \leq 6$, a totally segregated ∞ -bicyclic graph of order n does not exist.

Theorem 2.1

If G is a bicyclic graph with $\Delta(G)=k$, then $n_1(G) = n_3(G) + 2n_4(G) + \dots + (k-2)n_k(G) - 2$.

Proof. Let G be a bicyclic graph on n vertices. Then $n = n_1 + n_2 + \dots + n_k$. By first theorem of graph theory, $2(n+1) = n_1 + 2n_2 + \dots + kn_k$. Eliminating n and solving for n_1 , we get

$$n_1(G) = n_3(G) + 2n_4(G) + 3n_5(G) + \dots + (k-2)n_k(G) - 2. \blacksquare$$

Theorem 2.2

If G is a TSB graph with $\Delta(G)=k$, then $n_2(G) \leq (k-1)n_k + (k-2)n_{k-1} + \dots + 3n_4 + 2n_3 - 1$.

Proof. Let $G=(V, E)$ be a TSB graph with $\Delta(G)=k$. In G , if v is a vertex of degree 2 and $u, w \in N(v)$, then delete the vertex v and join the edge uw . After this vertex deletion and edge joining, degrees of u and w remain the same and number of vertices and edges are decreased by one. We continue this process till all vertices of degree 2 vanish. Let the resultant graph be G' . G' is again a bicyclic graph with $n_i(G') = n_i(G)$ for $i \neq 2$ and $n_2(G') = 0$. $|E(G')| = \sum_{i=3}^k n_i(G') + n_1(G') + 1 = \sum_{i=3}^k n_i(G) + n_1(G) + 1$. Since G is totally segregated $n_2(G) \leq |E(G')|$.

Then $n_2(G) \leq |E(G')| = \sum_{i=3}^k n_i(G) + n_1(G) + 1$. By theorem 2.1, we have

$$n_2(G) \leq (k-1)n_k + (k-2)n_{k-1} + \dots + 3n_4 + 2n_3 - 1. \blacksquare$$

Theorem 2.3

If $G = (V, E)$ is a totally segregated ∞ -bicyclic graph with $\Delta(G)=4$ and $n_4(G)=1$, then
 $n_3(G)-2 \leq n_2(G) \leq 2n_3(G)+2$.

Proof. Let G be a totally segregated ∞ -bicyclic graph. Let w be a common vertex of two cycles in G and $d_G(w)=4$. By theorem 2.2, $n_2(G) \leq 2n_3(G)+2$. In G , if v is a vertex of degree 2 and $u, w \in N(v)$, then delete the vertex v and join the edge uw . After this vertex deletion and edge joining, degrees of u and w remains the same and number of vertices and edges are decreased by one. We continue this process till all vertices of degree 2 vanish. Let the resultant graph be G' . G' is again a ∞ -bicyclic graph with $n_4(G')=1$, $n_3(G')=n_3(G)$, $n_2(G')=0$ and $n_1(G')=n_1(G)$. $|E(G')|=n_3(G') + n_1(G') + 2$ and G' has $n_3(G') + n_1(G') + 2 - (n_1(G') + 4) = n_3(G') - 2$ balanced edges. Hence $n_2(G) \geq n_3(G') - 2$, otherwise G will not be totally segregated graph. But $n_3(G')=n_3(G)$. Hence $n_2(G) \geq n_3(G) - 2$. ■

Remark 2.2

In a ∞ -bicyclic graph G on n vertices with $\Delta(G)=4$, $n_4(G)=1$.

$$n_2 \leq 2n_3 + 2 \Rightarrow n_3 \geq \frac{n-3}{4}, n_2 \leq 2n_3 + 2 \Rightarrow n_2 \leq \frac{n+1}{2}$$

$$n_2 \geq n_3 - 2 \Rightarrow n_3 \leq \frac{n+1}{3}, n_2 \geq n_3 - 2 \Rightarrow n_2 \geq \frac{n-5}{3}$$

Theorem 2.4

Let $k \geq 4$ be any positive integer, then for any integer x such that $k-2 \leq x \leq 2k+2$, there exists a totally segregated ∞ -bicyclic graph with $\Delta=4$, $n_4=1$, $n_2=x$, $n_3=k$, and $n_1=k$.

Proof. Consider a cycle C_{k-1} on $k-1$ ($k \geq 4$) vertices and the cycle C_3 . Let $w_1 \in V(C_{k-1})$ and $w_2 \in V(C_3)$. Identify w_1 and w_2 and the identified vertex is named as w . Add pendent edges to all vertices except w . Subdivide all edges of the cycles C_{k-1} and C_3 except the edges incident with w . Then subdivide $x - (k-2)$ edges among $k+4$ edges (k pendent edges and 4 edges incident with w). See Figure 3. The constructed graph is totally segregated ∞ -bicyclic graph G when $k=5$,
 $n_2(G)=x=k-2$. ■




Figure 3

Lemma 2.1

Let G be a totally segregated ∞ -bicyclic graph on n ($n \geq 9$) vertices and $\Delta(G)=k$ ($k > 3$). Then there exists ∞ -bicyclic graph, G^* on n vertices with $\Delta(G)=4$ and $n_4(G^*)=1$ satisfying the following conditions.

$$(i) \quad n_3(G^*) \geq \frac{n_2(G^*)-2}{2}$$

$$(ii) irr_t(G^*) < irr_t(G)$$

Proof. (i) Let G be a totally segregated ∞ -bicyclic graph on $n(n \geq 9)$ vertices and $\Delta(G) = k$ ($k \geq 4$). Let v_k, v_2 (if exists), v_1 be any vertex of degree $k, 2, 1$ respectively in G , and degree sequence of G is given as follows. $(k^{n_k}, (k-1)^{n_{k-1}}, \dots, 2^{n_2}, 1^{n_1})$. By Theorem 2.2 we have

$$n_2(G) \leq (k-1)n_k(G) + (k-2)n_{k-1}(G) + \dots + 3n_4(G) + 2n_3(G) - 1 \dots \dots \dots (3)$$

By Theorem 2.1 we have

$$n_1(G) = (k-2)n_k(G) + (k-3)n_{k-1}(G) + \dots + 2n_4(G) + n_3(G) - 2 \dots \dots \dots (4)$$

If $n_2(G) \neq 0$, do the branch-transformation on G from v_k to v_2 . If $n_2(G) = 0$, do the branch-transformation on G from v_k to v_1 . Repeat the branch-transformation on G till maximum degree of resulting graph G^* is 4 and $n_4(G^*) = 1$. If $N_{Br}(G)$ denotes the number of branch transformations that are done in G ,

$$N_{Br}(G) = (k-3)n_k(G) + (k-4)n_{k-1}(G) + \dots + 2n_5(G) + n_4(G) - 1 \dots \dots \dots (5)$$

We prove this lemma by the following two cases:

Case 1.

If $n_2(G) \leq N_{Br}(G)$. That is,

$$n_2(G) \leq (k-3)n_k(G) + (k-4)n_{k-1}(G) + \dots + 2n_5(G) + n_4(G) - 1 \dots \dots \dots (6)$$

By equation (6), in the process of branch transformation from v_k to v_2 , at some stage, number of vertices of degree two in the resulting bicyclic graph is zero. Then $n_2(G^*) = 0$ or 1.

ie $n_2(G^*) \leq 2n_3(G^*)$, That is $n_3(G^*) \geq \frac{n_2(G^*)}{2}$

Case 2.

If $n_2(G) > N_{Br}(G)$. That is, if

$$n_2(G) > (k-3)n_k(G) + (k-4)n_{k-1}(G) + \dots + 2n_5(G) + n_4(G) - 1 \dots \dots \dots (7)$$

When the process of branch transformation ends, equation (3)-(7) gives upper bound for $n_2(G^*)$

$$n_2(G^*) \leq 2n_k(G) + 2n_{k-1}(G) + \dots + 2n_5(G) + 2n_4(G) + 2n_3(G) \dots \dots \dots (8)$$

Number of branch transformations that is done in G and number of vertices of degree greater than or equal to three in G gives $n_3(G^*)$.

That is,

$$n_3(G^*) = (k-3)n_k(G) + (k-4)n_{k-1}(G) + \dots + 2n_5(G) + n_4(G) - 1 \\ + n_k(G) + n_{k-1}(G) + \dots + n_5(G) + n_4(G) + n_3(G)$$

$$n_3(G^*) = (k-2)n_k(G) + (k-3)n_{k-1}(G) + \dots + 3n_5(G) + 2n_4(G) + n_3(G) - 1 \dots \dots \dots (9)$$

$$\text{From equations (8) and (9), } n_2(G^*) \leq 2n_3(G^*) + 2$$

Hence, in both cases, $n_3(G^*) \geq \frac{n_2(G^*) - 2}{2}$.

(ii) We prove this result by the following two cases.

Case 1.

G'' is formed from G by branch transformation on G from v_k to v_2 . By this transformation only the number of vertices with degrees $k, k-1, 3$ and 2 are changed.

Degree sequence of G is $(k^{n_k}, (k-1)^{n_{k-1}}, \dots, 2^{n_2}, 1^{n_1})$

and degree sequence of G'' is $(k^{n_k-1}, (k-1)^{n_{k-1}+1}, \dots, 3^{n_3+1}, 2^{n_2-1}, 1^{n_1})$.

By equation (2) $\text{irr}_t(G'') = \text{irr}_t(G) - 2n_{k-1} - 2n_{k-2} - 2n_{k-3} - \dots - 2n_5 - 2n_4 - 2n_3 - 2$

Hence $\text{irr}_t(G'') < \text{irr}_t(G)$

Case 2.

G' is formed from G by branch transformation on G from v_k to v_1 . By this transformation only the number of vertices with degrees $k, k-1, 2$ and 1 are changed.

Degree sequence of G is $(k^{n_k}, (k-1)^{n_{k-1}}, \dots, 3^{n_3}, 2^{n_2}, 1^{n_1})$

Degree sequence of G' is $(k^{n_k-1}, (k-1)^{n_{k-1}+1}, \dots, 3^{n_3}, 2^{n_2+1}, 1^{n_1-1})$.

By equation (2) $\text{irr}_t(G') = \text{irr}_t(G) - 2n_{k-1} - 2n_{k-2} - 2n_{k-3} - \dots - 2n_5 - 2n_4 - 2n_3 - 2n_2 - 2$

Hence $\text{irr}_t(G') < \text{irr}_t(G)$

We keep on repeating this process and obtain a bicyclic graph G^* with $\Delta=4$ and $n_4(G^*)=1$.

Then $\text{irr}_t(G^*) \leq \text{irr}_t(G)$ ■

Lemma 2.2

If G^* is any ∞ - bicyclic graph on n vertices with $\Delta(G^*)=4$, $n_4(G^*)=1$ and $n_3(G^*) \geq \frac{n_2(G^*)-2}{2}$, there exists TSB graph G_1 , on n vertices with $\Delta(G_1) = 4$, $n_4(G_1) = 1$, $n_3(G_1) = \left\lceil \frac{n-3}{4} \right\rceil$ and $\text{irr}_t(G_1) \leq \text{irr}_t(G^*)$.

Proof. Let G^* be a ∞ - bicyclic graph on n vertices with $\Delta(G^*) = 4$, $n_4(G^*)=1$ and $n_3(G^*) \geq \frac{n_2(G^*)-2}{2}$. ie $n_3(G^*) \geq \frac{n-3}{4}$ by remark 2.2. i.e. $n_3(G^*) \geq \left\lceil \frac{n-3}{4} \right\rceil$. Let degree sequence of G^* be $(4^1, 3^{n_3}, 2^{n_2}, 1^{n_1})$ and v_3 a vertex of degree three in G^* . Let G' be the bicyclic graph obtained from G^* by branch transformation on G^* from v_3 to v_1 . Degree sequence of G' is $(4^1, 3^{n_3-1}, 2^{n_2+2}, 1^{n_1-1})$.

$\text{Irr}_t(G') = (n_3(G^*)-1)+2(n_2(G^*)+2)+3(n_1(G^*)-1)+(n_3(G^*)-1)(n_2(G^*)+2)+$

$2(n_3(G^*)-1)(n_1(G^*)-1)+(n_2(G^*)+2)(n_1(G^*)-1)$

$\text{Irr}_t(G') = \text{irr}_t(G^*) - 2n_2(G^*) - 4$

Hence $\text{irr}_t(G') < \text{irr}_t(G^*)$. Repeat the above branch transformation on G^* till number of vertices of degree 3 in the resulting graph G_1 is $\left\lceil \frac{n-3}{4} \right\rceil$. i.e. $n_3(G_1) = \left\lceil \frac{n-3}{4} \right\rceil$, $n_1(G_1) = \left\lceil \frac{n-3}{4} \right\rceil$,

$n_2(G_1) = n - 1 - 2\left\lceil \frac{n-3}{4} \right\rceil$. Here $\left\lceil \frac{n-3}{4} \right\rceil - 2 \leq n - 1 - 2\left\lceil \frac{n-3}{4} \right\rceil \leq 2\left\lceil \frac{n-3}{4} \right\rceil + 2$. By theorem 2.4, for any integer x s.t. $k-2 \leq x \leq 2k+2$, there exists totally segregated ∞ - bicyclic graph with $\Delta=4$, $n_4=1$, $n_2=x$, $n_3=k$, $n_1=k$. ie there exists totally segregated ∞ - bicyclic graph, G_1^* on n vertices such that $n_3(G_1^*) = \left\lceil \frac{n-3}{4} \right\rceil$,

$n_2(G_1^*) = n - 1 - 2\left\lceil \frac{n-3}{4} \right\rceil$, $n_1(G_1^*) = \left\lceil \frac{n-3}{4} \right\rceil$, and $\text{irr}_t(G_1^*) \leq \text{irr}_t(G^*)$. ■

Theorem 2.5

Let G be a totally segregated ∞ - bicyclic graph on n ($n \geq 9$) vertices. Then $\text{irr}_t(G) \geq 2n \left\lceil \frac{n-3}{4} \right\rceil - 2 \left\lceil \frac{n-3}{4} \right\rceil^2 - 2 \left\lceil \frac{n-3}{4} \right\rceil + 2n - 2$, and the equality holds if and only if $\Delta(G)=4$, $n_4(G)=1$, $n_3(G) = \left\lceil \frac{n-3}{4} \right\rceil$.

Corollary 2.1

Let G be a totally segregated ∞ - bicyclic graph on n vertices. Then,

1. $\text{irr}_t(G) \geq \frac{3n^2}{8} + \frac{3n}{2} - 2$, when $n=4k$, $k=3,4,\dots$. Equality holds if and only if $n_4(G)=1$, $n_3(G)=k$, $n_2(G)=2k-1$, $n_1(G)=k$.
2. $\text{irr}_t(G) \geq \frac{3n^2}{8} + \frac{5n}{4} - \frac{13}{8}$, when $n=1+4k$, $k=2,3,4,\dots$. Equality holds if and only if $n_4(G)=1$, $n_3(G)=k$, $n_2(G)=2k$, $n_1(G)=k$.
3. $\text{irr}_t(G) \geq \frac{3n^2}{8} + n - \frac{3}{2}$, When $n=2+4k$, $k=2,3,4,\dots$. Equality holds if and only if $n_4(G)=1$, $n_3(G)=k$, $n_2(G)=2k+1$, $n_1(G)=k$.
4. $\text{irr}_t(G) \geq \frac{3n^2}{8} + \frac{3n}{4} - \frac{13}{8}$, when $n=3+4k$, $k=2,3,4,\dots$. Equality holds if and only if $n_4(G)=1$, $n_3(G)=k$, $n_2(G)=2k+2$, $n_1(G)=k$ ■

Its degree sequence determines the total irregularity. Below is the degree sequence of a fully segregated ∞ -bicyclic graph for each n , with least Total Irregularity ($\min G_n$):

No. of vertices: n	Degree sequence of $\min G_n$
$n = 4k$, $k = 2, 3, \dots$	$(4^1, 3^k, 2^{2k-1}, 1^k)$
$n = 4k+1$, $k = 2, 3, \dots$	$(4^1, 3^k, 2^{2k}, 1^k)$
$n = 4k+2$, $k = 2, 3, \dots$	$(4^1, 3^k, 2^{2k+1}, 1^k)$
$n = 4k+3$, $k = 2, 3, \dots$	$(4^1, 3^k, 2^{2k+2}, 1^k)$

Remark 2.3

For $n \leq 6$ totally segregated ∞ - bicyclic graph on n vertices does not exist.

For $n=7,8$ totally segregated ∞ - bicyclic graph on n vertices with minimum total irregularity is presented in Figure 4.




Figure 4

References:

1. H. Abdo and D. Dimitrov. "The Total Irregularity of a Graph." Discrete Mathematics and Theoretical Computer Science.vol.16. no.1 (2015): 201-206.
2. H. Abdo and D. Dimitrov. "Non-Regular Graphs with Minimal Total Irregularity." Bulletin of the Australian Mathematical Society. vol.92. no.1 (2015): 1-10.
3. Y. Alavi, G. Chartrand, F. R. K. Chung, P. Erdos, R. L. Graham and O. R. Ollermann. "Highly Irregular Graphs." Journal of Graph Theory. vol.11. no.2 (1987) 235-249.
4. M. O. Albertson. "The Irregularity of a graph." Ars.Combin. vol. 46, (1987) 219-225.
5. F.K. Bell. "A Note on the Irregularity of Graphs." Linear Algebra and its Applications. vol.161. (1992): 45-54.
6. G. Chartrand, P.Erdos and O.R. Ollermann. "How to Define an Irregular Graph." College Math. J. vol.19. (1988): 36-42.
7. G.H. Fath-Tabar, I Gutman and R.Nasiri. "Extremely Irregular Trees." Bulletin T.CXLV de l'AcademieSerbe des Sciences et des Arts(Cl. Sci. Math. Natur.). vol.38.(2013).
8. I. Gutman, P. Hansen and H.Melot, "Variable Neighborhood Search for Extremal Graphs. 10. Comparison of Irregularity Indices for Chemical Trees." J. Chem. Inf. Model. vol. 45. no.2 (2005): 222-230. [doi:10.1021/ci0342775]
9. D.E. Jackson and R. Entringer, "Totally Segregated Graphs." Congress. Numer. vol. 55. (1986): 159-165.
10. O Ravi, R Senthil Kumar, A Hamari Choudhi, Weakly \sqsupseteq g-closed sets, Bulletin Of The International Mathematical Virtual Institute, 4, Vol. 4(2014), 1-9
11. O Ravi, R Senthil Kumar, Mildly Ig-closed sets, Journal of New Results in Science, Vol3,Issue 5 (2014) page 37-47
12. O Ravi, A senthil kumar R & Hamari Choudhi, Decompositions of \tilde{I} g-Continuity via Idealization, Journal of New Results in Science, Vol 7, Issue 3 (2014), Page 72-80.
13. O Ravi, A Pandi, R Senthil Kumar, A Muthulakshmi, Some decompositions of πg -continuity, International Journal of Mathematics and its Application, Vol 3 Issue 1 (2015) Page 149-154.
14. S. Tharmar and R. Senthil Kumar, Soft Locally Closed Sets in Soft Ideal Topological Spaces, Vol 10, issue XXIV(2016) Page No (1593-1600).
15. Y. Zhu, L. You and J. Yang. "The Minimal Total Irregularity of Some Classes of Graphs." Filomat. vol. 30. no.5. (2016): 1203-1211.
16. L.You, J.Yang, Y.Zhu, Z.You. "The maximal total irregularity of bicyclic graphs." Journal of Applied Mathematics. vol. 2014. (2014):1-9. [doi.org/10.1155/2014/785084]

*